

# Isolation and cross-species amplification of novel microsatellite loci in a charismatic marine mammal species, the northern elephant seal (*Mirounga angustirostris*)

Simona Sanvito · Alejandro Dueñas Meza ·  
Yolanda Schramm · Pedro Cruz Hernández ·  
Yareli Esquer Garrigos · Filippo Galimberti

Received: 31 July 2012 / Accepted: 6 August 2012  
© Springer Science+Business Media B.V. 2012

**Abstract** Due to its demographic history, the northern elephant seal is a charismatic species with a peculiar place in conservation biology. After having being almost exterminated by commercial sealing, and having being repeatedly declared extinct, the species has enjoyed a period of expansion at sustained rate. The low genetic variability produced by the bottleneck is apparently not affecting the viability of the species, but implies practical problems in the application of standard molecular ecology tools due to the lack of polymorphic markers. We developed novel microsatellite markers that, although showing a rather small variability, are a valuable addition to the molecular toolbox that can be used to study the species.

**Keywords** Microsatellites · Genetic variability · Northern elephant seal · *Mirounga angustirostris*

---

**Electronic supplementary material** The online version of this article (doi:10.1007/s12686-012-9741-3) contains supplementary material, which is available to authorized users.

---

S. Sanvito · F. Galimberti  
Elephant Seal Research Group, Milan, Italy

S. Sanvito (✉) · A. Dueñas Meza · Y. Schramm  
Facultad de Ciencias Marinas, Universidad Autónoma de Baja California, Km 103 Carretera Ensenada-Tijuana, Ensenada, Baja California, Mexico  
e-mail: simo\_esrg@eleseal.org

P. Cruz Hernández  
Centro de Investigaciones Biológicas del Noroeste, La Paz, Baja California Sur, Mexico

Y. Esquer Garrigos  
UMR BOREA IRD 207, Muséum National d'Histoire Naturelle, 43 rue Cuvier, 75005 Paris, France

The northern elephant seal (*Mirounga angustirostris*; NES) is an important species for conservation biology theories because of its peculiar demographic history, and the genetic consequences of it. The species suffered a significant population bottleneck due to commercial sealing, was repeatedly declared extinct, and was actually reduced to a very small number of individuals (20–100; Weber et al. 2000), concentrated on a single island, Isla Guadalupe (Mexico). The species shows a very low genetic variability in all markers that have been tested, including allozymes (Bonnell and Selander 1974), mtDNA (Weber et al. 2000), minisatellites (Lehman et al. 1993) and MHC loci (Weber et al. 2004). Notwithstanding this sustained depletion of genetic variability, after Mexico and USA government adopted strict protection rules the species enjoyed an impressive recovery (Bartholomew and Hubbs 1960), growing at a sustained rate, and greatly increasing its geographic range (Stewart et al. 1994; Carretta et al. 2011). All together, it seems that the lack of variability is not affecting the viability of the species, at least in the medium term (Weber et al. 2004). The lack of variability has an important negative practical implication: it is difficult to apply to NES the molecular ecology methods that are normally used to study population structure, kinship, paternity, etc., because of the lack of suitable polymorphic markers. In this note we present the development of 9 new polymorphic microsatellite markers that are a valuable addition to the molecular toolbox that can be used to study this species.

We collected skin samples in the Islas San Benito (Baja California, Mexico), taking by surprise small (1–2 g) pieces of inter-digital membrane of the rear flippers of live NES, and storing them in 2.0 ml screw cap tubes with 95 % ethanol kept at ambient temperature (Fabiani et al. 2004). DNA was extracted from skin samples using the

DNEasy Blood and Tissue kit (Qiagen), following the manufacturer mouse-tail protocol. Extracted DNA was checked for degradation by electrophoresis in 1.0 % agarose gels and by NanoDrop ND-1000 (ThermoScientific)

measurements of absorbance (DNA concentration between 100 and 300  $\mu\text{g}/\mu\text{l}$ ).

We used 6 male and 6 female samples to construct four microsatellite enriched libraries using biotinylated probes

**Table 1** Characteristics of the novel loci

Locus/GenBank	Repeat motif	Primers	PCR	Conc.	MgCl <sub>2</sub>	T <sub>a</sub>	N	N <sub>a</sub>	Size
Mang01 JQ714261	(CT) <sub>2</sub> GT(CT) <sub>4</sub> ...(CT) <sub>10</sub>	GCCTTTGGTTAGGTATCCAG CACTCTGAATACTGTAGCCTTG	Good	0.300 0.300	1.50	55	22	2	212–216
Mang03 JQ714262	(TA) <sub>5</sub>	GTGGAAAGAGCCAAGATG CTCAGCATAGTATCCTCTAGTTCC	Good	0.300 0.075	1.50	55	22	1	147
Mang04 JQ714263	(GA) <sub>9</sub>	GAGCTCTAGGTTATGATTTG CTGCTGCTCTACCAGC	Good	0.300 0.075	1.50	55	22	1	105
Mang05 JQ714264	(GT) <sub>5</sub>	GTTGTCGTGAGGACTGATG GAGATGAGCCTTAAAGAATGG	Good	0.300 0.075	1.50	55	22	1	112
Mang06 JQ714265	(GT) <sub>9</sub> ...(GA) <sub>14</sub>	CTATCACGGAGATGGGTG CTCAGGAAACCTTCATTGC	Good	0.300 0.075	1.50	55	22	2	159–161
Mang09 JQ714266	(TATC) <sub>6</sub> CATC(TATC) <sub>7</sub> ATC(TATC) <sub>3</sub> ...(TATC) <sub>12</sub>	GGAAGGAGGTGCTATTACTCTC CTCATCTCTTGAGGCATCC	Good	0.300 0.075	1.50	52	22	6	230–242
Mang14 JQ714267	(CT) <sub>3</sub> GT(CT) <sub>3</sub>	GAGCCTCCTCTGTAATGG GATTCGTGACCAGAAAATC	Good	0.300 0.300	1.50	55	22	1	129
Mang16 JQ714268	(CT) <sub>7</sub>	CAGAACATCAAACCAAGTGAG GGTTCAGTGTCTGCCTTC	Good	0.300 0.075	1.50	59	22	1	227
Mang17 JQ714269	(CA) <sub>6</sub> TG(CA) <sub>5</sub>	GAGTGTACCCCTTCCTCG GCAAAATGCTGTGTATGAGC	Good	0.300 0.075	1.50	55	22	1	153
Mang21 JQ714270	(GT) <sub>5</sub> CTCT(GT) <sub>3</sub>	GATAACTTCTGGGGTGGG GCTGAGAAAAATACTGTAAGATTC	Good	0.300 0.075	1.50	55	22	1	132
Mang23 JQ714271	(CA) <sub>11</sub>	CAGTGACTTCCCCCTCC GATCACAGGACAGCCTTCAG	Scarce	0.300 0.075	1.25	57	22	1	136
Mang27 JQ714272	(GT) <sub>2</sub> GA(GT) <sub>16</sub>	GGAAATGGTATTGTAGTTATGTAGG CTCCCCCTTCTGCATC	Good	0.300 0.075	1.50	55	22	2	109–111
Mang33 JQ714273	(CT) <sub>3</sub> T(CT) <sub>6</sub>	CCTGGTGGCTCTTGATT TGACACATTACAAAATACTCCA	Good	0.300 0.075	1.50	55	22	1	273
Mang34 JQ714274	(GT) <sub>7</sub> GCA(TG) <sub>6</sub> CA(TG) <sub>4</sub>	GCTGATGGACTGGCATTTTA GTGCTCGCCTCCTCTCCT	Scarce	0.300 0.075	1.70	60	22	3	196–200
Mang35 JQ714275	(CA) <sub>14</sub>	ATTGGTTTCTTGATTATGC ATGCCCGTATCTATTCCT	Good	0.300 0.075	1.50	53	22	2	245–248
Mang36 JQ714276	(CCAT) <sub>3</sub> ...(CCAT) <sub>4</sub> ...(CCAT) <sub>2</sub> ...(CCAT) <sub>5</sub>	GGGGACACAAGCACAAC CTCAAAGGATGGATAGATAAGC	Good	0.300 0.075	1.50	55	22	2	339–343
Mang37 JQ714277	(TC) <sub>4</sub>	GAGCCCCGCATCAGG TTTATTTATTTAGAGAGTTCGTG	Good	0.300 0.075	1.50	57	22	1	104
Mang38 JQ714278	(GATA) <sub>4</sub> GATTA(GATA) <sub>13</sub>	GGGGACAGCACAAGGAAG GAAGGAATGGGAAGCCTA	Good	0.300 0.075	1.50	56	21	1	217
Mang41 JQ714279	(GATA) <sub>3</sub> ...(GATA) <sub>6</sub> GAT(GATA) <sub>2</sub> GAT(GATA) <sub>3</sub> ...(GATA) <sub>3</sub> ...(GATA) <sub>3</sub> GAT(GATA) <sub>10</sub>	GCCTTTCCTTCTTTCC GTCTCCATAACTGCCTGA	Good	0.300 0.075	1.50	53	22	1	276
Mang43 JQ714280	(GATA) <sub>2</sub> GAT(GATA) <sub>11</sub>	ACAGGATAGGGAATGGTGA GGGAAAGAGGATTGTTC	Good	0.300 0.075	1.50	55	21	3	238–246
Mang44 JQ714281	(GATA) <sub>14</sub> ...(GATA) <sub>2</sub> GAT(GATA) <sub>2</sub>	CATCTTACCAGGAGACAG GAGACAAGGATAGGTCA	Good	0.300 0.075	1.50	55	22	3	176–188
Mang48 JQ714282	(GT) <sub>16</sub> (GA) <sub>10</sub>	AGCCTGTAGCCCTTGT GCACCTTCTGTGTGAG	Good	0.300 0.075	1.50	55	22	1	270

GenBank = GenBank accession number; Primers = primer sequence, forward above and reverse below; PCR = quality of the amplification; Conc. = primer concentration, forward above and reverse below ( $\mu\text{M}$ ); MgCl<sub>2</sub> = magnesium concentration (mM); T<sub>a</sub> = annealing temperature ( $^{\circ}\text{C}$ ); N = number of typed individuals; N<sub>a</sub> = number of alleles; Size = allele size range in base pairs excluding the M13 universal tag

(CT)<sub>15</sub>, (GT)<sub>15</sub>, (CTGT)<sub>6</sub> and (GATA)<sub>10</sub>, following the protocol of Glenn and Schable (2005). The sequencing of 160 positive clones showed the presence of 89 microsatellite loci (success rate = 55.6 %), 56 % dinucleotides and 30 % tetranucleotides. We designed primers for 48 of these loci (Primer3, Rozen, and Skaletsky 2000).

For the PCR amplification and genotyping we used the universal tag approach of Schuelke (2000). We used the following PCR mix (15 µl final volume): 1X PCR buffer (Promega), 1.0–1.9 (see Table 1 and electronic supplement) mM MgCl<sub>2</sub>, 200 µM dNTP Mix, 0.3 µM forward primer, 0.3 or 0.075 µM (see Table 1) reverse primer plus M13 universal tag (5' TGTAACGACGGCCAGT 3'), 0.3 µM M13 universal tag plus fluorescent dye (Hex, Fam, Ned), and 0.02 units of GoFlexi Taq DNA polymerase (Promega). We used the following PCR program: 5' at 95 °C; 35 cycles of 20" at 95 °C, 20" at the optimized annealing temperature (see Table 1) and 30" at 72 °C; 8 cycles of 20" at 95 °C, 20" at 47 °C and 30" at 72 °C; 10' at 72 °C; hold at 4 °C. PCR products were resolved on a 3730XL Automated Sequencer (Applied Biosystems), and genotyped using GeneMarker 1.85 (SoftGenetics). We genotyped 22 NES individuals. We used the same protocol to optimize PCR conditions and carry out the genotyping for 5 individuals of each of the following pinniped species: southern elephant seal (*M. leonina*, ML), harbor seal (*Phoca vitulina*, PV), California sea lion (*Zalophus californianus*, ZC), and Guadalupe fur seal (*Arctocephalus twonsoni*, AT). Samples of these species were collected during previous research projects carried out by the Authors.

We calculated statistics of microsatellite variation and exclusion probabilities using custom scripts, we tested

the loci for Hardy–Weinberg equilibrium in GenePop4 (Rousset 2008), we verified the presence of null alleles in Microchecker (Van Oosterhout et al. 2004), and we calculated maximum likelihood estimates of genotyping error and null alleles frequency in ML-Null (Kalinowski and Taper 2006).

We were able to optimize PCR conditions and carry out the genotyping for 22 loci (Table 1). We found a low variability, with a mean number of alleles per locus of 1.7 (SD = 1.2, range = 1–6). Only 9 loci (40.9 %) were polymorphic, with a mean number of alleles per locus of 2.8 (SD = 1.3; loci statistics in Table 2). All polymorphic loci were in Hardy–Weinberg equilibrium (all loci test, Fisher's method: *P* = 0.89; Table 2). We were able to cross-amplify most loci in all the species tested (AT 86.4 %, ML 100 %, PV 95.5 %, ZC 81.8 %; Table 1 of the Electronic Supplement). We obtained for all the four species a greater allele range than for NES.

The results of our study were mixed. Although we were able to develop new microsatellite markers for the species, their variability was rather limited, and not greater than the variability observed in microsatellites developed in other species and cross-amplified in NES (unpublished data). Anyway, we think that these new microsatellite loci are a valuable addition to the molecular toolbox that can be used to study northern elephant seals, in particular if combined with the cross-amplification of microsatellites developed for other pinniped species. On a more general ground, our study confirms that NES have an unusually low variability also in microsatellite loci. Observed variability statistics are lower than the ones calculated in most seals and sea lions, and are similar to the ones found in two heavily

**Table 2** Statistics of the polymorphic loci

Locus	H <sub>o</sub>	H <sub>e</sub>	F <sub>is</sub>	I	N <sub>e</sub>	PIC	PE <sub>1</sub>	PE <sub>2</sub>	PE <sub>3</sub>	PID	PID <sub>sibs</sub>	β	Null	P(HW)
Mang01	0.545	0.495	-0.103	0.676	1.936	0.367	0.183	0.117	0.276	0.371	0.604	0.000	0.000	0.682
Mang06	0.091	0.088	-0.024	0.184	1.094	0.083	0.042	0.004	0.078	0.818	0.916	0.000	0.000	1.000
Mang09	0.636	0.734	0.133	1.419	3.538	0.669	0.475	0.303	0.656	0.103	0.424	0.000	0.006	0.236
Mang27	0.273	0.305	0.104	0.474	1.424	0.253	0.127	0.044	0.207	0.502	0.736	0.000	0.027	0.538
Mang34	0.591	0.559	-0.057	0.928	2.204	0.486	0.291	0.149	0.441	0.231	0.543	0.000	0.000	0.202
Mang35	0.364	0.359	-0.012	0.536	1.541	0.290	0.145	0.062	0.230	0.450	0.695	0.000	0.000	1.000
Mang36	0.273	0.240	-0.132	0.398	1.307	0.208	0.104	0.028	0.176	0.577	0.785	0.000	0.000	1.000
Mang43	0.190	0.182	-0.005	0.383	1.217	0.169	0.086	0.014	0.159	0.650	0.833	0.045	0.000	1.000
Mang44	0.773	0.625	-0.237	1.019	2.570	0.541	0.331	0.186	0.482	0.195	0.500	0.000	0.000	0.216
Mean	0.415	0.399	-0.037	0.669	1.870	0.341						0.005	0.004	
SD	0.230	0.217	0.114	0.387	0.794	0.192						0.015	0.009	
Cumulative							0.881	0.635	0.972	0.000	0.021			

H<sub>o</sub> = observed heterozygosity; H<sub>e</sub> = gene diversity (expected heterozygosity); F<sub>is</sub> = inbreeding coefficient; I = Shannon index of information; N<sub>e</sub> = number of effective alleles; PIC = polymorphism information content; PE<sub>1</sub> = one parent probability of exclusion; PE<sub>2</sub> = missing parent probability of exclusion; PE<sub>3</sub> = Both parents probability of exclusion; PID = probability of identity; PID<sub>sibs</sub> = probability of identity among sibs; β = maximum likelihood estimate of genotypic error; Null = maximum likelihood estimate of null allele frequency; P(HW) = exact probability of deviation from Hardy–Weinberg equilibrium

bottlenecked phocid species, the Mediterranean monk seal (Pastor et al. 2004) and the Hawaiian monk seal (Schultz et al. 2010) that, contrary to NES, are demographically challenged.

**Acknowledgments** For logistic support during the field work we would like to thank the Cooperativa Pescadores Nacionales de Abulon of Isla de Cedros. We also would like to thank the Secretaria de Medio Ambiente y Recursos Naturales for granting the research licenses. We would like to thank Professor Faustino Camarena Rosales for providing to us laboratory space.

## References

- Bartholomew GA, Hubbs CL (1960) Population growth and seasonal movements of the northern elephant seal, *Mirounga angustirostris*. *Mammalia* 24:313–324
- Bonnell ML, Selander RK (1974) Elephant seals: genetic variation and near extinction. *Science* 184:908–909
- Carretta JV, Forney KA, Muto MM, Barlow J, Baker JD, Hanson B, Lowry MS (2011) U.S. Pacific marine mammal stock assessments: 2010. NOAA Technical Memorandum NMFS NMFS NOAA-TM-NMFS-SWFSC-476. U.S. Department of Commerce, National Oceanic and Atmospheric Administration, La Jolla
- Fabiani A, Galimberti F, Sanvito S, Hoelzel AR (2004) Extreme polygyny among southern elephant seals on Sea Lion Island, Falkland Islands. *Behav Ecol* 15:961–969
- Glenn TC, Schable NA (2005) Isolating microsatellite DNA loci. In: Zimmer E, Roalson E (eds) *Methods in enzymology, Molecular evolution: Producing the Biochemical Data Part B*, vol 395, Academic Press, San Diego, pp 202–222
- Kalinowski S, Taper M (2006) Maximum likelihood estimation of the frequency of null alleles at microsatellite loci. *Conserv Genet* 7:991–995
- Lehman N, Wayne RK, Stewart BS (1993) Comparative levels of genetic variability in harbour seals and elephant seals. *Symp Zool Soc Lond* 66:49
- Pastor T, Garza JC, Allen P, Amos W, Aguilar A (2004) Low genetic variability in the highly endangered Mediterranean monk seal. *J Heredity* 95:291–300
- Rousset F (2008) GENEPOP'007: a complete re-implementation of the GENEPOP software for Windows and Linux. *Mol Ecol Resour* 8:103–106
- Rozen S, Skaletsky HJ (2000) Primer3 on the WWW for general users and for biologist programmers. In: Krawetz S, Misener S (eds) *Bioinformatics methods and protocols: methods in molecular biology*. Humana Press, Totowa, pp 365–386
- Schuelke M (2000) An economic method for the fluorescent labeling of PCR fragments. *Nat Biotechnol* 18:233–234
- Schultz JK, Marshall AJ, Pfunder M (2010) Genome-wide loss of diversity in the critically endangered Hawaiian monk seal. *Diversity* 2:863–880
- Stewart BS, Yochem PK, Huber HR, DeLong RL, Jameson RJ, Sydeman WJ, Allen SG, Le Boeuf BJ (1994) History and present status of the northern elephant seal population. In: Le Boeuf BJ, Laws RM (eds) *Elephant seals. Population ecology, behavior and physiology*. University of California Press, Berkeley, pp 29–48
- Van Oosterhout C, Hutchinson WF, Wills DPM, Shipley P (2004) MICRO-CHECKER: software for identifying and correcting genotyping errors in microsatellite data. *Mol Ecol Notes* 4: 535–538
- Weber DS, Stewart BS, Garza JC, Lehman N (2000) An empirical genetic assessment of the severity of the northern elephant seal population bottleneck. *Curr Biol* 10:1287–1290
- Weber DS, Stewart BS, Schienman J, Lehman N (2004) Major histocompatibility complex variation at three class II loci in the northern elephant seal. *Mol Ecol* 13:711–718